

## Real time aurora oval forecasting - SvalTrackII

F. Sigernes<sup>(1)</sup>, M. Dyrland<sup>(1)</sup>, P. Brekke<sup>(2)</sup>, E. K. Gjengedal<sup>(3)</sup>, S. Chernouss<sup>(4)</sup>,  
D. A. Lorentzen<sup>(1)</sup>, K. Oksavik<sup>(1)</sup>, C. S. Deehr<sup>(5)</sup>

1. The University Centre in Svalbard (UNIS), N-9171 Longyearbyen, Norway.

2. Norwegian Space Centre, Oslo, Norway.

3. Storm Weather Center, Bergen, Norway.

4. Polar Geophysical Institute, Murmansk Region, Apatity, Russia.

5. Geophysical Institute, University of Alaska, Fairbanks, USA.

(\*) Email: Fred.Sigernes@unis.no

Recibido / Received: 30/01/2011. Aceptado / Accepted: 30/08/2011.

### ABSTRACT:

A method to forecast, up to one hour in the future, the size and location of the aurora oval is described. The work is based on a mathematical description of the aurora oval coupled to predicted values of the planetary  $K_p$  index. As a result, a real time animation of the oval mapped onto the Earth's surface is created. The night- and dayside are visualized together with the location of the twilight zone as Earth rotates under the aurora oval.

**Keywords:** Space Physics, Aurora Ovals, Forecast.

### REFERENCIAS Y ENLACES / REFERENCES AND LINKS

- [1]. Y. I. Feldstein, "Auroral oval", *J. Geophys. Res.* **78**, 1210-1213 (1973).
- [2]. R. H. Holzworth, C. I. Meng, "Mathematical representation of the auroral oval", *Geophys. Res. Lett.* **2**, 377-380 (1975).
- [3]. M. S. Gussenhoven, D. A. Hardy, N. Heinemann, "Systematics of the equatorward diffuse auroral boundary", *J. Geophys. Res.* **88**, 5692-5708 (1983).
- [4]. K. A. Costello, *Moving the Rice MSFM into a Real-Time Forecast Mode Using Solar Wind Driven Forecast Models*, Ph.D. dissertation, Rice University, Houston, TX (1997).
- [5]. G. V. Starkov, "Mathematical model of the auroral boundaries", *Geomagnet. Aeron.* **34**, 331-336 (1994).
- [6]. G. V. Starkov, "Statistical dependences between the magnetic activity indices", *Geomagnet. Aeron.* **34**, 101-103 (1994).
- [7]. D. A. Vallado, P. Crawford, R. Hujsak, T. S. Kelso, "Revisiting Spacetrack Report #3", American Institute of Aeronautics and Astronautics (AIAA), Report No AIAA 2006-6753, 1-88 (2006).

### 1. Introduction

The impact areas on the Earth of energetic particles from the Sun causing circular belts of auroral emissions around each geomagnetic magnetic pole are known as the aurora ovals. The location and size of these ovals have been studied extensively during the last decades [1-3].

The size and location of the auroral oval was found to be directly related to the planetary index of geomagnetic activity,  $K_p$ . This index is

produced by the Helmholtz Centre in Potsdam. Because the production of this index involves worldwide data collection and interpretation in order to continue a record begun in 1936, it is not useful for forecasting, and an intermediate method of scaling  $K_p$  from one or a small number of mid-latitude stations has evolved. The  $K_p$  index derived in this way must be referred to as the estimated  $K_p$  index, and it is this index we use in the present work.

The advent of stationary spacecraft approximately one hour upstream in the solar wind between the Sun and the Earth has led to studies of the relationship between the structure of the solar wind and the resulting auroral and geomagnetic disturbances [4]. Because the input data come from a satellite approximately one hour upstream in the solar wind, the resultant forecast is a relatively short-term forecast. This can be a very useful quantity both for auroral observing and for the purpose of conducting experiments that are dependent on the location and intensity of the aurora.

During the International Geophysical Year (IGY 1957-1958), the boundaries of auroral occurrence were determined from all-sky camera studies for a wide variety of activity levels. The resulting figure was made up from a statistical study of auroral occurrence showing the poleward and equatorward boundaries of the 75% occurrence probability [1]. The relationship between the morphology of the auroral oval and the level of geomagnetic activity allows us to develop a means of locating the aurora, independent of the vagaries of auroral observations. In addition, we may predict its location up to an hour in the future.

The goal of this paper is therefore to produce in real time, a forecast, up to one hour in the future, of the location and size of the aurora oval mapped on to the Earth's surface that includes observing conditions from a selected geographical site. The solar elevation and the cloud cover are the main obstacles for an aurora watcher at ground level. To the authors' knowledge, no such display of information exists.

Below is a mathematical presentation of the aurora oval and the geographical mapping that is used to visualize the results.

## 2. A mathematical representation of the aurora ovals

The model used to calculate the boundaries of the aurora oval is from work done by Starkov [5]. He compiled simple formulas of the poleward, the equatorward and the diffuse

aurora boundaries as a function of magnetic activity.

The magnetic input parameter is the AL index. This index describes the polar or planetary magnetic disturbances that occur during auroras. The range of the index is in the order of  $\pm 800$  nT. In terms of  $K_p$  index it is [6]

$$AL = 18 - 12.3K_p + 27.2K_p^2 - 2K_p^3. \quad (1)$$

The value of the  $K_p$  index varies from 0 to 9 with 0 being very quiet and 5 or more indicating geomagnetic storm conditions. In detail,  $K_p$  represents a 3 hour weighted average from a global network of magnetometers measuring the maximum deviation in the horizontal component of the Earth's magnetic field.

In coordinates of corrected geomagnetic colatitude,  $\theta$ , the boundaries of the oval are expressed as

$$\begin{aligned} \theta = A_0 + A_1 \cos[15(t + \alpha_1)] \\ + A_2 \cos[15(2t + \alpha_2)] \quad (2) \\ + A_3 \cos[15(3t + \alpha_3)] \end{aligned}$$

where  $A_i$  for  $i \in [0 \dots 3]$  are amplitudes in units of degrees of latitude,  $t$  is the local time in decimal hours, and  $\alpha_i$  for  $i \in [1 \dots 3]$  are phases also given in units of hours. Equation (2) is valid for the poleward, the equatorward and the diffuse aurora boundaries. The challenge now is to find the coefficients of Eq. (2). Starkov [5] does this using a new third order polynomial for both the  $A_i$  and  $\alpha_i$  coefficients

$$\begin{aligned} A_i \text{ or } \alpha_i = b_0 + b_1 \log_{10}|AL| + b_2 \log_{10}^2|AL| \\ + b_3 \log_{10}^3|AL| \quad (3) \end{aligned}$$

In other words, for each coefficient  $A_i$  or  $\alpha_i$  there is a set of  $b_i$  for  $i \in [0 \dots 3]$  values. Table I lists all the values. Starting with the  $K_p$  index, Eq. (1) is used to obtain the AL index. Next, Table I is used to obtain the correct set of  $b_i$  values to calculate the  $A_i$  or  $\alpha_i$  coefficients by Eq. (3). Finally, the auroral boundaries are obtained using Eq. (2). Note that the above procedure works for  $K_p$  indices that are floating point numbers. The ovals are expanding smoothly as the  $K_p$  index increases.

TABLE 1  
Expansion coefficients for auroral boundaries by Starkov [5] used in Eq. (3).

	$A_0$	$A_1$	$A_2$	$A_3$	$\alpha_1$	$\alpha_2$	$\alpha_3$
Poleward boundary of the auroral oval							
$b_0$	-0.07	-10.06	-4.44	-3.77	-6.61	6.37	-4.48
$b_1$	24.54	19.83	7.47	7.90	10.17	-1.10	10.16
$b_2$	-12.53	-9.33	-3.01	-4.73	-5.80	0.34	-5.87
$b_3$	2.15	1.24	0.25	0.91	1.19	-0.38	0.98
Equatorward boundary of the auroral oval							
$b_0$	1.61	-9.59	-12.07	-6.56	-2.22	-23.98	-20.07
$b_1$	23.21	17.78	17.49	11.44	1.50	42.79	36.67
$b_2$	-10.97	-7.20	-7.96	-6.73	-0.58	-26.96	-24.20
$b_3$	2.03	0.96	1.15	1.31	0.08	5.56	5.11
Equatorward boundary of the diffuse aurora							
$b_0$	3.44	-2.41	-0.74	-2.12	-1.68	8.69	8.61
$b_1$	29.77	7.89	3.94	3.24	-2.48	-20.73	-5.34
$b_2$	-16.38	-4.32	-3.09	-1.67	1.58	13.03	-1.36
$b_3$	3.35	0.87	0.72	0.31	-0.28	-2.14	0.76

### 3. Geographical transform

The ovals calculated in the previous section are centred on the magnetic poles. The Cartesian components in the graph are related to the polar magnetic coordinates by

$$\begin{aligned} x_m &= \sin\theta\cos\phi; & y_m &= \sin\theta\sin\phi; \\ z_m &= \cos\theta, \end{aligned} \tag{4}$$

where  $\theta$  and  $\phi$  are the magnetic latitude and longitude, respectively. In local magnetic time it becomes

$$\phi = \frac{2\pi t}{24} + \Delta\phi(t). \tag{5}$$

$\Delta\phi(t)$  is the longitudinal difference between the sub-solar point and the magnetic poles at time  $t$  (hours). It is important in order to make sure that the ovals are oriented correctly with magnetic noon pointing towards the Sun as Earth rotates around its own axis. The transformation to geographic coordinates is

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \cos\phi'_0\cos\lambda & -\sin\phi'_0 & \cos\phi'_0\sin\lambda \\ \sin\phi'_0\cos\lambda & \cos\phi'_0 & \sin\phi'_0\sin\lambda \\ -\sin\lambda & 0 & \cos\lambda \end{bmatrix} \begin{bmatrix} x_m \\ y_m \\ z_m \end{bmatrix}. \tag{6}$$

The invariant magnetic north pole is located in geographic coordinates at latitude  $\theta'_0=82.42^\circ N$

and longitude  $\phi'_0=-82.86^\circ E$ .  $\lambda = \pi/2 - \theta'_0$  is the latitudinal difference between the geographic and the north magnetic pole. Finally, the geographic latitude and longitude of the ovals are given as

$$\begin{aligned} \theta &= \frac{\pi}{2} - \cos^{-1}(z), \\ \phi' &= \begin{cases} \psi & \forall x > 0 \\ \psi + \pi & x < 0 \end{cases}, \\ \psi &= \tan^{-1}\left(\frac{y}{x}\right) \end{aligned} \tag{7}$$

Note that the procedure is identical for the south magnetic pole if we assume that the  $K_p$  index is the same.

### 4. Visualization

The ovals are visualized with a stand alone 32-bit executable Windows program called SvalTrackII. The program is written in Borland's Delphi - Pascal. It displays interactively mapping data in real time onto a three-dimensional spherical Globe representing the Earth. The twilight zone, night- and dayside of the Earth are projected with grades of shade on the Globe as a function of time. The 3D globe can be rotated and zoomed to display a close-up of any region of the Earth.

Both the aurora Borealis and the aurora Australis ovals are projected as polygons onto the globe with an angular resolution of  $1.5^\circ$ . The equatorward boundary of the diffuse aurora is added as a polygonal line. The local position of the aurora observer is added as a point with corresponding state information of the Moon and the Sun. In addition, the circle of  $\sim 4.5^\circ$  around the observer represents a  $160^\circ$  field of sky view, assuming that the auroral emissions peaks at an altitude of  $\sim 110$  km. The program also maps the position of space objects. The orbits are calculated by the use of the Simplified General Perturbations model 4 for near and deep space objects (SGP4 / SDP4) [7]. The model input is compatible with the Two-Line Element sets (TLE data) from NORAD (North American Aerospace Defense Command).

Figure 1 shows all the features. A textured map is used to visualize the Earth on 24<sup>th</sup> of December 2009. The twilight zone crosses most of Norway, and North America is on the night side. The observer is located on the island




Fig. 1: SvalTrackII screen dump: (1) equatorward boundary of the diffuse aurora, (2) aurora oval  $K_p=3$ , (3) magnetic north pole, (4) field of view aurora observer, (5) observer location, (6) Moon and Sun information at local site, (7) position of the Rapideye 1 satellite. Time: 09:00 UT on 24<sup>th</sup> December 2009.

of Svalbard in Longyearbyen, Norway ( $78.2^\circ N$ ,  $16.0^\circ E$ ). The  $K_p = 3$  oval is typically broader on the night- vs. the dayside, with magnetic noon or cusp located over the site at  $\sim 08:50$  UT. The Moon is below the horizon, indicating favorable conditions to view the aurora. The satellite named Rapideye 1 is located at coordinates ( $73.4^\circ N$ ,  $62.9^\circ E$ ) at an altitude of 631.5 km. The corresponding azimuth- and elevation angle for the observer are  $86.9^\circ$  to the East and  $19.3^\circ$  above the horizon, respectively. Several satellites may be tracked simultaneously.

Figure 2 shows different sized aurora ovals as a function of  $K_p$  index and time of day for 24<sup>th</sup> December 2009. As expected, the impact area or ovals are wider and more asymmetric with increasing  $K_p$  index. Note that the island of Svalbard is uniquely located. It is possible to view both the day- and nightside aurora from Svalbard.

The only input to the above visualization is time, TLE data and the estimated  $K_p$  index. The program is set up to update itself based on the predicted  $K_p$  index provided by the Space Weather Prediction Centre at the National Oceanic and Atmospheric Administration (NOAA). The index is predicted every 15 minutes by the use of the Costello Geomagnetic Activity Index model [4], which takes into account the most recent response of solar wind parameters (neural network algorithm).

The net result is an aurora oval forecast up to approximately one hour in advance of real time installed to run automatically at the Kjell Henriksen Observatory (KHO) on Svalbard, Norway.

## 5. Concluding remark

A method to mathematically calculate the size and location of the aurora oval onto a solar illuminated Earth globe is presented as function of estimated  $K_p$  index and time. The system may be used as a forecast when the  $K_p$  value is estimated from satellite data one hour upstream in the solar wind.




Fig. 2: Animated aurora ovals as a function of  $K_p$  index [0...8] and time for 24<sup>th</sup> December 2009.

### Acknowledgements

We wish to thank NOAA - Space Weather Prediction Centre for allowing us to download the predicted value of the  $K_p$  index. The work is financially supported by The Research Council of Norway through the project: Norwegian and Russian Upper Atmosphere Co-operation On Svalbard 2 # 196173/S30 and the Nordic Council of Ministers: Arctic Cooperation Programme # A10162.